



Classification of algebras of level two in the variety of nilpotent algebras and Leibniz algebras

James Francese^a, Abror Khudoyberdiyev^{b,*}, Bennett Rennier^c,
Anastasia Voloshinov^d

^a Fullerton College, Fullerton, CA, 92832, USA

^b National University of Uzbekistan, Institute of Mathematics, Tashkent, 100125, Uzbekistan

^c University of Oklahoma, Norman, OK, 73019, USA

^d Pomona College, Claremont, CA, 91711, USA

ARTICLE INFO

Article history:

Received 4 April 2018

Accepted 27 August 2018

Available online xxxx

MSC:

14D06

14L30

17A32

Keywords:

Leibniz algebra

Nilpotent algebra

Degenerations

Level of algebra

ABSTRACT

This paper is devoted to the description of complex finite-dimensional algebras of level two. We obtain the classification of algebras of level two in the variety of Leibniz algebras. It is shown that, up to isomorphism, there exist three Leibniz algebras of level two, one of which is solvable, and two of which are nilpotent. Moreover, we describe all algebras of level two in the variety of nilpotent algebras.

© 2018 Elsevier B.V. All rights reserved.

1. Introduction and background

The theory of deformations and degenerations of algebras has its origins in certain formal relations between physical theories and has become a lively subject in algebraic and differential geometry, as well as noncommutative and nonassociative algebra. It was a very influential precept of Niels Bohr that a new physical theory, which is supposed to ontologically overlap with a previously accepted theory, should somehow yield the old theory as a limiting or special case [1]. This is a statement of his “correspondence principle”, which is realized in quantum mechanics via the limit of the Moyal bracket:

$$[f, g] = \{f, g\} + \mathcal{O}(\hbar^2)$$

as $\hbar \rightarrow 0$. Here the bracket $\{, \} = \partial^i \partial_i - \partial_i \partial^i$ is the Poisson bracket of classical mechanics, being a sum of commutators for first-order differential operators. In this way classical mechanics emerges, as a limiting case, from quantum mechanics for small values of \hbar . This realization motivated the modern deformation theory of algebras, which originated with Gerstenhaber [2] and others. This theory has powerful applications in the classification of algebraic varieties and the quantization theory of Poisson manifolds, where the physical meaning of deformation remains especially explicit [3,4]. There

* Corresponding author.

E-mail addresses: jamesafrancese@gmail.com (J. Francese), khabror@mail.ru (A. Khudoyberdiyev), bennett@brennier.com (B. Rennier), nastia13@gmail.com (A. Voloshinov).

we have a manifold with tangent space V equipped with a Poisson bivector $p \in V \wedge V^*$, and an algebra of observables $f, g \in C^\infty(V)$. We then define a product of these observables as a power series in a parameter \hbar :

$$(f \star g)(p) := f \cdot g + \hbar p_j^i \frac{\partial f}{\partial x^i} \frac{\partial g}{\partial x_j} + \frac{1}{2} \hbar^2 p_l^k p_j^i \frac{\partial^2 f}{\partial x^k \partial x^i} \frac{\partial^2 g}{\partial x_l \partial x_j} + \dots$$

where we have made use of the summation convention. Defining the bracket

$$[f, g] := -i\hbar^{-1}(f \star g - g \star f),$$

in the ordinary quantum mechanical case for a single state (i.e. where all differential operators are first-order) we recover the above limit, relating the Moyal to the Poisson bracket. The full higher-order approach is currently being used in the study of formal quantum field theory [5,6], a version of Hilbert’s 6th problem [7].

For this paper, we consider the case of a finite-dimensional algebra over a closed field k , with $k = \mathbf{C}$ being of special focus. In general, an algebra over k can be considered as an element $\mu \in \text{Hom}(V \otimes V, V)$, where V is an n -dimensional k -vector space. Thus, in the purely abstract deformation theory we consider algebras $\mu, \mu_t \in V \otimes k[[t]]$ related by a formal power series:

$$\mu_t = \mu + \sum_{i=1}^{\infty} t^i \varphi_i \quad \text{where } \varphi_i \in \text{Hom}(V \otimes V, V)$$

so that in the linear case $\mu_{t,1} = \mu + t\varphi_1$ we study algebras with multiplication differing by a 2-coboundary.¹ Kodaira and Spencer gave the original idea of infinitesimal deformations for complex analytic manifolds [8]. Most notably, they proved that infinitesimal deformations can be parametrized by a related cohomology group. In fact, cohomology detects deformation at all orders, and it is therefore unsurprising that one can develop the deformation theory in any abelian category [9].

Prior to the development of deformation theory, it had already been realized that the space and time symmetries of Newtonian mechanics were recovered in the $c \rightarrow \infty$ limit of special relativity, where c is the speed of light. In that case the Lie algebra of the Poincare group degenerates to the Lie algebra of the Galilean group, an observation first made by İnönü and Wigner in [10]. This is a geometric process, and can be defined purely in terms of the Zariski topology on $\text{Hom}(V \otimes V, V)$.

In the finite-dimensional case over a field of characteristic zero, such degenerations can be described directly in terms of the singular limit of a linear group action. Let $\text{Alg}_n(k)$ be the variety of n -dimensional algebras over k , and let $\lambda, \mu \in \text{Alg}_n(k)$. Define an action on $\text{Alg}_n(k)$ by means of

$$(g \star \mu)(x, y) := g(\lambda(g^{-1}(x), g^{-1}(y))) \quad \text{where } g \in \text{GL}_n(k), \quad x, y \in A$$

which just represents a change of basis for A as an algebra. Thus the orbit of the algebra (A, λ) under this action is given by

$$\text{Orb}(A) := \{L \in \text{Alg}_n(k) \mid L \simeq A\}.$$

Definition 1. An algebra (A, λ) is said to degenerate to the algebra (A, μ) if $\text{Orb}(A, \mu) \subseteq \overline{\text{Orb}(A, \lambda)}$. We write $\lambda \rightarrow \mu$ to denote this degeneration.

In the case $k = \mathbf{C}$, we have that $\lambda \rightarrow \mu$ if and only if there is a $g_t \in \text{GL}_n(\mathbf{C}(t))$ such that $\forall x, y \in A$,

$$\mu(x, y) = \lim_{t \rightarrow 0} g_t(\lambda(g_t^{-1}(x), g_t^{-1}(y))).$$

We call a degeneration $\lambda \rightarrow \mu$ *trivial* if $(A, \lambda) \simeq (A, \mu)$, and *direct* if it is non-trivial, and there is no algebra (A, ν) such that $\lambda \rightarrow \nu \rightarrow \mu$. If $\lambda \rightarrow \mu$, then λ is a non-trivial deformation of μ , thus it is common to pass from the degeneration theory to the deformation theory.

It is clear that every non-abelian algebra in $\text{Alg}_n(\mathbf{C})$ degenerates non-trivially to the abelian algebra ab_n , but of course not all such degenerations will be direct; the distance of an algebra from ab_n , in terms of the degeneration theory, is given by its *level*.

Definition 2. The level of an algebra λ is the maximum length of a chain of direct degenerations to ab_n . We denote the level of an algebra by $\text{lev}_n(\lambda)$.

Concerning algebras of level one, we have the following result proved by Khudoyberdiyev and Omirov [11].

Theorem 1. Let A be an algebra of level one. Then A is isomorphic to one of the following pairwise non-isomorphic algebras:

$$\begin{aligned} p_n^- : & \quad e_1 e_i = e_i, \quad e_i e_1 = -e_i, \quad 2 \leq i \leq n; \\ n_3^- \oplus \text{ab}_{n-3} : & \quad e_1 e_2 = e_3, \quad e_2 e_1 = -e_3; \\ \lambda_2 \oplus \text{ab}_{n-2} : & \quad e_1 e_1 = e_2; \\ \nu_n(\alpha) : & \quad e_1 e_1 = e_1, \quad e_1 e_i = \alpha e_i, \quad e_i e_1 = (1 - \alpha) e_i, \quad 2 \leq i \leq n. \end{aligned}$$

The level two case, within the varieties of Lie, Jordan, and associative algebras, has been resolved by Khudoyberdiyev in [12]. In particular, that paper provides the following theorem.

¹ An abstract ‘‘Poisson bracket’’ on an algebra is always available in the case of a linear deformation, since we can define $\{x, y\} := \frac{1}{2}(\mu_{t,1}(x, y) - \mu_{t,1}(y, x))$.

Theorem 2. Let G be a Lie algebra of level two. Then G is isomorphic to one of the following pairwise non-isomorphic algebras:

$$\begin{aligned} n_{5,1} \oplus ab_{n-5} &: e_1e_3 = e_5, & e_2e_4 = e_5, & 2 \leq i \leq n; \\ n_{5,2} \oplus ab_{n-5} &: e_1e_2 = e_4, & e_1e_3 = e_5; \\ r_2 \oplus ab_{n-2} &: e_1e_1 = e_2; \\ g_{n,1}(\alpha) &: e_1e_2 = \alpha e_2, & e_1e_i = e_i, & 3 \leq i \leq n, \alpha \in \mathbf{C} \setminus \{0, 1\}; \\ g_{n,2} &: e_1e_2 = e_2 + e_3, & e_1e_i, & 3 \leq i \leq n. \end{aligned}$$

It is still desirable to obtain a complete classification of level two algebras. One step is to ask about the existence of level two algebras in other varieties. Since $n_{5,1}$ and $n_{5,2}$ are Lie, we may well ask if there are any non-Lie Leibniz algebras of level two.

Definition 3 ([13]). A (right) Leibniz algebra is a non-associative algebra such that for all $x, y, z \in L$, the following identity holds:

$$x(yz) = (xy)z - (xz)y.$$

This is a natural generalization of Lie algebras, in that an antisymmetric Leibniz algebra is Lie. Note that a left Leibniz algebra is defined by identity $(xy)z = x(yz) - y(xz)$.

Degenerations of Lie and Leibniz algebras were the subject of numerous papers, see for instance [14–18] and references given therein, and their research continues actively. In particular, in [19,20] some irreducible components of Leibniz algebras are found.

In this paper, we extend Theorem 2 to identify all non-Lie Leibniz algebras of level two; we find that two of these are nilpotent and one is solvable. We then proceed to classify all n -dimensional nilpotent algebras of level two, and find that these are all Leibniz.

2. Main results

Our first main result is the classification of Leibniz algebras of level two.

Theorem 3. Let L be a n -dimensional non-Lie Leibniz algebra of level two. Then L is isomorphic one of the following three algebras:

$$\begin{aligned} L_4(\alpha) \oplus ab_{n-3} &: e_1e_1 = e_3, & e_2e_1 = e_3, & e_2e_2 = \alpha e_3; \\ L_5 \oplus ab_{n-3} &: e_1e_1 = e_3, & e_1e_2 = e_3, & e_2e_1 = e_3; \\ r_n &: e_i e_1 = e_i, & 2 \leq i \leq n. \end{aligned}$$

Together with the nilpotent Lie algebras of level two identified in [12], our other main result identifies these four algebras as the only nilpotent algebras of level two.

Theorem 4. Any finite-dimensional nilpotent algebra of level two is isomorphic to one of the following algebras:

$$\begin{aligned} n_{5,1} \oplus ab_{n-5} &: e_1e_3 = e_5, & e_2e_1 = e_5; \\ n_{5,2} \oplus ab_{n-5} &: e_1e_2 = e_4, & e_1e_3 = e_5; \\ L_4(\alpha) \oplus ab_{n-3} &: e_1e_1 = e_3, & e_2e_2 = \alpha e_3, & e_1e_2 = e_3; \\ L_5 \oplus ab_{n-3} &: e_1e_1 = e_3, & e_1e_2 = e_3, & e_2e_1 = e_3. \end{aligned}$$

It is now natural to ask if any algebra of level two is a direct sum of two level one algebras. The following examples give us a negative answer to this question.

Example 1. The algebras

$$\begin{aligned} n_3^- \oplus \lambda_2 &= \{x_1, x_2, x_3, x_4, x_5\} : x_1x_1 = x_2, & x_3x_4 = x_5, & x_4x_3 = -x_5; \\ \lambda_2 \oplus \lambda_2 &= \{x_1, x_2, x_3, x_4\} : x_1x_1 = x_2, & x_3x_3 = x_4; \\ \lambda_2 \oplus p_n^- &= \{x_1, x_2, x_3, x_4, \dots, x_n\} : x_1x_1 = x_2, & x_ix_3 = x_i, & x_3x_i = -x_i, \quad 4 \leq i \leq n. \end{aligned}$$

via the family of matrices

$$\left\{ \begin{array}{l} g_t^{-1}(x_1) = t(x_1 + x_3), \\ g_t^{-1}(x_2) = \frac{t}{2}(x_1 + x_4), \\ g_t^{-1}(x_3) = t^2x_2, \\ g_t^{-1}(x_4) = t^2x_1, \\ g_t^{-1}(x_5) = t(x_5 + x_2), \end{array} \right\} \left\{ \begin{array}{l} g_t^{-1}(x_1) = tx_1, \\ g_t^{-1}(x_2) = t(x_1 + x_3), \\ g_t^{-1}(x_3) = t^2x_2, \\ g_t^{-1}(x_4) = t(x_2 + x_4), \end{array} \right\} \left\{ \begin{array}{l} g_t^{-1}(x_1) = t(x_1 + x_4), \\ g_t^{-1}(x_2) = t(\frac{1}{2}x_1 + x_3), \\ g_t^{-1}(x_3) = t^2x_2, \\ g_t^{-1}(x_4) = t(x_4 + \frac{1}{2}x_2), \\ g_t^{-1}(x_i) = tx_i, \end{array} \right\} \quad 5 \leq i \leq n,$$

degenerate to the algebras $L_4(\frac{1}{4}) \oplus ab_2$, $L_5 \oplus ab_1$ and $L_4(\frac{1}{4}) \oplus ab_{n-3}$ respectively.

Since the algebras $L_4(\alpha)$ and L_5 are not algebras of level one, we deduce that the level of the algebras $n_3^- \oplus \lambda_2$, $\lambda_2 \oplus \lambda_2$ and $\lambda_2 \oplus p_n^-$ must be greater than two.

Now let L be a n -dimensional complex algebra and $\{e_1, e_2, \dots, e_n\}$ be a basis of L . The multiplication on the algebra L is defined by the products of the basis elements; namely, by the products

$$e_i e_j = \sum_{k=1}^n \gamma_{ij}^k e_k,$$

where γ_{ij}^k are the structural constants.

We first prove a very useful lemma, which will allow us to immediately conclude a degeneration to either $L_4(\alpha)$ or L_5 based on a multiplication table of a certain form.

Lemma 1. *Suppose L is an n -dimensional algebra and let $\{e_1, e_2, \dots, e_n\}$ be a basis of L . If there exist distinct i, j, k such that*

$$(\gamma_{i,i}^k, \gamma_{i,j}^k, \gamma_{j,i}^k, \gamma_{j,j}^k) \notin \{(0, \beta, -\beta, 0), (\delta, \beta, \beta, \frac{\beta^2}{\delta})\} \text{ where } \delta \neq 0,$$

then $L \rightarrow L_4(\alpha)$ or $L \rightarrow L_5$.

Proof. Without loss of generality, we may assume $i = 1, j = 2$, and $k = 3$. We see that if we take the degeneration

$$g_t(e_1) = t^{-1}e_1, \quad g_t(e_2) = t^{-1}e_2, \quad g_t(e_i) = t^{-2}e_i \quad 3 \leq i \leq n,$$

then we have following nontrivial products

$$\begin{aligned} e_1 e_1 &= \gamma_{1,1}^3 e_3 + \sum_{s=4}^n \gamma_{1,1}^s e_s, & e_1 e_2 &= \gamma_{1,2}^3 e_3 + \sum_{s=4}^n \gamma_{1,2}^s e_s, \\ e_2 e_1 &= \gamma_{2,1}^3 e_3 + \sum_{s=4}^n \gamma_{2,1}^s e_s, & e_2 e_2 &= \gamma_{2,2}^3 e_3 + \sum_{s=4}^n \gamma_{2,2}^s e_s. \end{aligned}$$

Furthermore, if we take the additional degeneration

$$g_t(e_1) = t^{-1}e_1, \quad g_t(e_2) = t^{-1}e_2, \quad g_t(e_3) = t^{-2}e_3, \quad g_t(e_i) = t^{-1}e_i \quad 4 \leq i \leq n,$$

then we have an algebra with the following multiplication

$$e_1 e_1 = \gamma_{1,1}^3 e_3, \quad e_1 e_2 = \gamma_{1,2}^3 e_3, \quad e_2 e_1 = \gamma_{2,1}^3 e_3, \quad e_2 e_2 = \gamma_{2,2}^3 e_3.$$

We see that this algebra is nilpotent and also non-Lie as $(\gamma_{1,1}^3, \gamma_{1,2}^3, \gamma_{2,1}^3, \gamma_{2,2}^3) \neq (0, \beta, -\beta, 0)$. Moreover, since $(\gamma_{1,1}^3, \gamma_{1,2}^3, \gamma_{2,1}^3, \gamma_{2,2}^3) \neq (\delta, \beta, \beta, \frac{\beta^2}{\delta})$, we conclude that L is not isomorphic to the algebra λ_2 . Due to the classification of three dimensional nilpotent Leibniz algebras [21], we conclude that this algebra is isomorphic to either $L_4(\alpha)$ or L_5 . \square

2.1. Classification of algebras of level two in the variety of Leibniz algebras

Let L be a Leibniz algebra and let $x \in L$. We define $\varphi_x : L \rightarrow L$ to be the linear operator where $\varphi_x(y) = yx + xy$. We see that by applying the Leibniz identity we get the following two equations:

$$\begin{aligned} z\varphi_x(y) &= z(yx + xy) = z(yx) + z(xy) = (zy)x - (zx)y + (zx)y - (zy)x = 0, \\ z(xx) &= (zx)x - (zx)x = 0. \end{aligned}$$

This proves that both $\varphi_x(y)$ and xx are in the right annihilator for any $x \in L$.

In this section we will examine the matrix representation of φ_x on a case-by-case basis in order to prove [Theorem 3](#).

Proposition 1. *Let L be a n -dimensional non-Lie Leibniz algebra which is not of level one (i.e. $L \not\cong \lambda_2$). Then L degenerates to one of the following three algebras:*

$$\begin{aligned} L_4(\alpha) \oplus \mathfrak{ab}_{n-3} : & \quad e_1 e_1 = e_3, & \quad e_2 e_1 = e_3, & \quad e_2 e_2 = \alpha e_3; \\ L_5 \oplus \mathfrak{ab}_{n-3} : & \quad e_1 e_1 = e_3, & \quad e_1 e_2 = e_3, & \quad e_2 e_1 = e_3; \\ r_n : & \quad e_i e_1 = e_i, & \quad 2 \leq i \leq n. \end{aligned}$$

Proof. Since L is a non-Lie Leibniz algebra, we know that there exists an element $x \in L$ such that $xx \neq 0$. Suppose that $xx = \alpha x$ for some constant α . Then, by the Leibniz identity, we have that

$$\alpha^2 x = \alpha xx = x(\alpha x) = x(xx) = (xx)x - (xx)x = 0$$

which means that $\alpha = 0$. This is a contradiction, though, as $xx \neq 0$. Thus, it must be that xx is linearly independent from x . Using this, we can form a basis $\{e_1, e_2, \dots, e_n\}$, where $e_1 = x$ and $e_2 = xx$.

We define the linear operator $\varphi = \varphi_x$ and let $(\alpha_{i,j})$ be its matrix form. Thus, we have that

$$\varphi = \begin{bmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & \alpha_{1,4} & \dots & \alpha_{1,n} \\ 2 & \alpha_{2,2} & \alpha_{2,3} & \alpha_{2,4} & \dots & \alpha_{2,n} \\ 0 & \alpha_{3,2} & \alpha_{3,3} & \alpha_{3,4} & \dots & \alpha_{3,n} \\ 0 & \alpha_{4,2} & \alpha_{4,3} & \alpha_{4,4} & \dots & \alpha_{4,n} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & \alpha_{n,2} & \alpha_{n,3} & \alpha_{n,4} & \dots & \alpha_{n,n} \end{bmatrix}$$

Suppose that $\alpha_{j,k} \neq 0$ for some $1, j, k$ distinct and $k \geq 3$. This means that we have the following products

$$e_j e_1 = e_2, \quad e_j e_j = \gamma_{j,j}^k e_k + \sum_{s=1, s \neq k}^n \gamma_{j,j}^s e_s,$$

$$e_1 e_j = \gamma_{1,j}^k e_k + \sum_{s=1, s \neq k}^n \gamma_{1,j}^s e_s, \quad e_j e_1 = (\alpha_{j,k} - \gamma_{1,j}^k) e_k + \sum_{s=1, s \neq k}^n \gamma_{j,1}^s e_s.$$

Since $\gamma_{1,1}^k = 0$ and $\gamma_{1,j}^k \neq -(\alpha_{j,k} - \gamma_{1,j}^k)$, we can apply Lemma 1 on the indices $1, j, k$ to see that $L \rightarrow L_4(\alpha)$ or L_5 . Now we focus the case where $\alpha_{j,k} = 0$ for $1, j, k$ distinct and $k \geq 3$. This gives us the following matrix representation of φ :

$$\varphi = \begin{bmatrix} 0 & \alpha_{1,2} & \alpha_{1,3} & \alpha_{1,4} & \dots & \alpha_{1,n} \\ 2 & \alpha_{2,2} & \alpha_{2,3} & \alpha_{2,4} & \dots & \alpha_{2,n} \\ 0 & 0 & \alpha_{3,3} & 0 & \dots & 0 \\ 0 & 0 & 0 & \alpha_{4,4} & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \alpha_{n,n} \end{bmatrix}$$

Suppose that $\alpha_{2,i} \neq 0$ for some $i \neq 1, 2$. Without loss of generality, let $i = 3$. Then if we take the change of basis $e'_3 = e_1 - \frac{2}{\alpha_{2,3}} e_3$, we have that

$$\varphi(e'_3) = \varphi(e_1 - \frac{2}{\alpha_{2,3}} e_3) = \varphi(e_1) - \frac{2}{\alpha_{2,3}} \varphi(e_3) = 2e_2 - \frac{2}{\alpha_{2,3}} (\alpha_{1,3} e_1 + \alpha_{2,3} e_2 + \alpha_{3,3} e_3)$$

$$= -\frac{2\alpha_{1,3}}{\alpha_{2,3}} e_1 - \frac{2\alpha_{3,3}}{\alpha_{2,3}} e_3 = -\frac{2\alpha_{1,3}}{\alpha_{2,3}} e_1 - \alpha_{3,3} (e_1 - e'_3) = -\left(\frac{2\alpha_{1,3}}{\alpha_{2,3}} - \alpha_{3,3}\right) e_1 + \alpha_{3,3} e'_3.$$

Thus, we may assume that $\alpha_{2,3} = 0$. Since $i = 3$ was arbitrary, we can therefore assume that $\alpha_{2,i} = 0$ for $3 \leq i \leq n$. This means that the matrix representation of φ has the form:

$$\varphi = \begin{bmatrix} 0 & \alpha_{12} & \alpha_{13} & \alpha_{14} & \dots & \alpha_{1n} \\ 2 & \alpha_{22} & 0 & 0 & \dots & 0 \\ 0 & 0 & \alpha_{33} & 0 & \dots & 0 \\ 0 & 0 & 0 & \alpha_{44} & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & \alpha_{nn} \end{bmatrix}$$

Now suppose there exists $j \geq 3$ such that $\alpha_{2,2} \neq \alpha_{j,j}$. Then we see that if we take the basis change $e'_j = e_2 + e_j$, we have that

$$\varphi(e'_j) = \varphi(e_2) + \varphi(e_j) = (\alpha_{1,2} + \alpha_{1,j})e_1 + \alpha_{2,2}e_2 + \alpha_{j,j}e_j$$

$$= (\alpha_{1,2} + \alpha_{1,j})e_1 + (\alpha_{2,2} - \alpha_{j,j})e_2 + \alpha_{j,j}e'_j$$

By reapplying our previous argument, we see that it must be that either $L \rightarrow L_4(\alpha)$, L_5 or $\alpha_{2,2} - \alpha_{j,j} = 0$. Thus, we have that $\alpha = \alpha_{2,2} = \alpha_{i,i}$ for $3 \leq i \leq n$.

Now suppose that $\alpha \neq 0$. We can then take the basis change $e'_i = \varphi(e_i)$ for $2 \leq i \leq n$. This means that $e_2, e_3, \dots, e_n \in \text{Ann}_R(L)$ and thus that $\varphi(e_i) = e_i e_1$ for $2 \leq i \leq n$. Moreover, if $\alpha_{1,i} \neq 0$ for some $2 \leq i \leq n$, then we would have that

$$\alpha_{1,i} e_1 = \varphi(e_i) - \alpha e_i \in \text{Ann}_R(L)$$

which is a contradiction to the fact that $e_1 e_1 = e_2$. Therefore, we have an algebra L with the following multiplication table:

$$e_1 e_1 = e_2, \quad e_i e_1 = \alpha e_i, \quad (\alpha \neq 0), \quad 2 \leq i \leq n.$$

If we then take the basis transformation $e'_1 = \frac{1}{\alpha}(e_1 - \frac{1}{\alpha}e_2)$, we see that this is exactly the algebra r_n .

Suppose then that $\alpha = 0$. Again, we see that if $\alpha_{1,i} \neq 0$ for some $2 \leq i \leq n$, then we would have that $\alpha_{1,i}e_1 = \varphi(e_i) \in \text{Ann}_R(L)$, which is a contradiction. This means that φ has the following matrix representation:

$$\varphi = \begin{bmatrix} 0 & 0 & 0 & 0 & \dots & 0 \\ 2 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \dots & 0 \end{bmatrix}$$

We now consider following products

$$e_2e_2 = 0, \quad e_3e_2 = 0, \quad e_2e_3 = \gamma_{2,3}^2e_2 + \gamma_{2,3}^3e_3 + \sum_{s=4}^n \gamma_{2,3}^s e_s, \quad e_3e_3 = \gamma_{3,3}^2e_2 + \gamma_{3,3}^3e_3 + \sum_{s=4}^n \gamma_{3,3}^s e_s.$$

If there exist $\gamma_{2,3}^s \neq 0$, for some $4 \leq s \leq n$, then by Lemma 1, we have that $L \rightarrow L_4(\alpha), L_5$. Thus, we assume that $\gamma_{2,3}^s = 0$ for $4 \leq s \leq n$. Therefore, $e_2e_3 = \gamma_{2,3}^2e_2 + \gamma_{2,3}^3e_3$. If $\gamma_{2,3}^3 \neq 0$, then we have that $\gamma_{2,3}^3e_3 = e_2e_3 - \gamma_{2,3}^2e_2 \in \text{Ann}_R(L)$. This means that $e_2e_3 = 0$, a contradiction as $\gamma_{2,3}^3 \neq 0$. Hence, $e_2e_3 = \gamma_{2,3}^2e_2$.

Assume that $\gamma_{2,3}^2 \neq 0$. If we take the change of basis $e'_2 = e_2 - Ae_1$, then we have that

$$e'_2e'_2 = (e_2 - Ae_1)(e_2 - Ae_1) = A^2e_1e_1 = A^2e_2 = A^3e_1 + A^2e'_2,$$

$$e'_2e_3 = (e_2 - Ae_1)e_3 = \gamma_{2,3}^2e_2 - Ae_1e_3 = A(\gamma_{2,3}^2 - \gamma_{1,3}^1 - A\gamma_{1,3}^2)e_1 + (\gamma_{2,3}^2 - A\gamma_{1,3}^2)e'_2 - A \sum_{s=3}^n \gamma_{1,3}^s e_s,$$

$$e_3e'_2 = e_3(e_2 - Ae_1) = Ae_1e_3 = A(\gamma_{1,3}^1 + A\gamma_{1,3}^2)e_1 + A\gamma_{1,3}^2e'_2 + A \sum_{s=3}^n \gamma_{1,3}^s e_s,$$

$$e_3e_3 = \gamma_{3,3}^1e_1 + \gamma_{3,3}^2e_1 + \sum_{s=3}^n \gamma_{3,3}^s e_s.$$

Since $\gamma_{2,3}^2 \neq 0$, we see that we can always choose A so that $\gamma_{2,3}^2 - \gamma_{1,3}^1 - A\gamma_{1,3}^2 \neq \gamma_{1,3}^1 + A\gamma_{1,3}^2$. Hence, using Lemma 1 we obtain that $L \rightarrow L_4(\alpha), L_5$. Thus, it must be that $\gamma_{2,3}^2 = 0$, which means that $e_2e_3 = 0$. Since e_3 was arbitrary, we may assume that $e_2e_i = 0$ for $i \neq 1, 2$ and thus we have that $e_2 \in \text{Ann}(L)$.

Therefore, we have the following multiplication table

$$e_1e_1 = e_2, \quad e_2e_i = e_ie_2 = 0, \quad 1 \leq i \leq n,$$

$$e_1e_i = -e_ie_1 = \sum_{s=1}^n \gamma_{1,i}^s e_s, \quad 3 \leq i \leq n,$$

$$e_ie_j = \sum_{s=1}^n \gamma_{i,j}^s e_s, \quad 3 \leq i, j \leq n.$$

Using the Leibniz identity

$$0 = (e_1e_1)e_i - (e_1e_i)e_1 + e_1(e_ie_1) = e_2e_i - \left(\sum_{s=1}^n \gamma_{1,i}^s e_s \right) e_1 - e_1 \left(\sum_{s=1}^n \gamma_{1,i}^s e_s \right) = -2\gamma_{1,i}^1 e_2$$

we obtain that $\gamma_{1,i}^1 = 0$ for $3 \leq i \leq n$. Furthermore, if $\gamma_{1,i}^2 \neq 0$ or $\gamma_{i,i}^2 \neq 0$ for some $3 \leq i \leq n$, then by Lemma 1, we have that $L \rightarrow L_4(\alpha), L_5$. Therefore, we may assume $\gamma_{1,i}^2 = 0$ and $\gamma_{i,i}^2 = 0$ for any $3 \leq i \leq n$.

Moreover, if $\gamma_{i,j}^2 \neq 0$ for some i, j then taking the basis change $e'_1 = e_1 + e_j$, we obtain that

$$e_1e_1 = e_2 + \sum_{i=3}^n (*)e_i, \quad e_ie_1 = \gamma_{i,j}^2e_2 + \sum_{i=3}^n (*)e_i,$$

$$e_ie_1 = \gamma_{j,i}^2e_2 + \sum_{i=3}^n (*)e_i, \quad e_ie_i = \sum_{i=3}^n (*)e_i.$$

If we apply Lemma 1 on the indices $1, i, 2$, we see that $L \rightarrow L_4(\alpha), L_5$. Therefore we can suppose $\gamma_{i,j}^2 = 0$ for any $3 \leq i, j \leq n$.

By applying the Leibniz identity once again we see that

$$0 = (e_i e_j) e_1 - (e_i e_1) e_j - e_i (e_j e_1) = \gamma_{ij}^1 e_2 + \sum_{i=1, i \neq 2}^n (*) e_i,$$

which means that $\gamma_{ij}^1 = 0$ for $3 \leq i, j \leq n$.

Therefore, we obtain that any Leibniz algebras L either degenerates to $L_4(\alpha)$, L_5 or r_n or is a decomposed algebra with ideals $M_1 = \{e_1, e_2\}$ and $M_2 = \{e_3, e_4, \dots, e_n\}$. If M_2 were trivial, then our only nontrivial multiplication would be $e_1 e_1 = e_2$. This is a contradiction, as we assumed that $L \not\cong \lambda_2$. Thus, M_2 is not trivial, and therefore M_2 degenerates to an algebra of level one: λ_2, n_3^- or p_n^- . This means that L degenerates to $\lambda_2 \oplus \lambda_2, \lambda_2 \oplus n_3^-$, or $\lambda_2 \oplus p_n^-$. By Example 1, we conclude that L degenerates to either $L_4(\alpha)$ or L_5 . \square

Theorem 5. Let L be an n -dimensional Leibniz algebra of level two. Then L is isomorphic to one of the following pairwise non-isomorphic algebras:

$$L_4(\alpha) \oplus \text{ab}_{n-3}, \quad L_5 \oplus \text{ab}_{n-3}, \quad r_n, \quad \alpha \in \mathbb{C}$$

Proof. Due to Theorem 3, it is sufficient to prove that these algebras do not degenerate to each other. To facilitate this, we compute the dimensions of right annihilator and derivations of these algebras and we call upon the following table:

$$\begin{aligned} \dim \text{Ann}_R(L_5) &= n - 2, & \dim \text{Der}(L_4(0)) &= n^2 - 3n + 4, \\ \dim \text{Ann}_R(L_4(\alpha)) &= n - 2, \quad \alpha \neq 0, & \dim \text{Der}(r_n) &= (n - 1)^2 = n^2 - 2n + 1, \\ \dim \text{Ann}_R(r_n) &= n - 1. \end{aligned}$$

We first note that $L_4(\alpha)$ and L_5 cannot degenerate to r_n , as $L_4(\alpha)$ and L_5 are nilpotent and r_n is not. We also see that r_n does not degenerate to $L_4(\alpha)$ ($\alpha \neq 0$) or L_5 , as dimension of right annihilator of r_n is more than dimensions of right annihilators of $L_4(\alpha)$ ($\alpha \neq 0$) and L_5 . Additionally, since for $n \geq 4$, the dimension of derivations of r_n is more than the dimension of derivations of $L_4(0)$, we have that $r_n \not\rightarrow L_4(0)$. Lastly, we see that $L_4(\alpha) \not\rightarrow L_5$ and that $L_5 \not\rightarrow L_4(\alpha)$ by the following paper [14]. \square

Remark 1. We note that in the context of left Leibniz algebras, the following algebra replaces the algebra r_n as a Leibniz algebra of level two:

$$\ell_n : \quad e_1 e_i = e_i, \quad 2 \leq i \leq n.$$

2.2. Nilpotent algebras of level two

Working in the variety of n -dimensional nilpotent algebras $\text{Nil}_n(\mathbb{C})$ will allow us to exclude certain products from our multiplication tables, in particular all products of the form $xy = x$.

Theorem 6. Any n -dimensional ($n \geq 5$) nilpotent algebra of level two is isomorphic to one of the following algebras:

$$\begin{aligned} n_{5,1} : \quad & e_1 e_2 = e_5, \quad e_3 e_4 = e_5, \quad e_2 e_1 = -e_5, \quad e_4 e_3 = -e_5; \\ n_{5,2} : \quad & e_1 e_2 = e_4, \quad e_1 e_3 = e_5, \quad e_2 e_1 = -e_4, \quad e_3 e_1 = -e_5; \\ L_4(\alpha) : \quad & e_1 e_1 = e_3, \quad e_2 e_2 = \alpha e_3, \quad e_1 e_2 = e_3; \\ L_5 : \quad & e_1 e_1 = e_3, \quad e_1 e_2 = e_3, \quad e_2 e_1 = e_3. \end{aligned}$$

Proof. Our overall strategy is to look separately at antisymmetric and non-antisymmetric cases, and then at the way products fall into the square of the algebra A^2 .

Case 1. First, we assume that $A \in \text{Nil}_n(\mathbb{C})$ is non-antisymmetric.

Case 1.1. Assume that $\dim(A^2) = 1$, then we assume that $A^2 = \{e_n\}$ and have the following multiplication

$$A : \begin{cases} e_1 e_1 = e_n, & e_i e_j = \alpha_{i,j} e_n, & 2 \leq i, j \leq n - 1, \\ e_1 e_j = \alpha_{1,j} e_n, & e_j e_1 = \alpha_{j,1} e_n, & 2 \leq j \leq n - 1. \end{cases}$$

If there exist i such that $|\alpha_{1i} - \alpha_{i1}| + |\alpha_{ii} - \alpha_{1i}\alpha_{i1}| \neq 0$, then by Lemma 1 we obtain that L degenerates to $L_4(\alpha)$ or L_5 . Now let $|\alpha_{1i} - \alpha_{i1}| + |\alpha_{ii} - \alpha_{1i}\alpha_{i1}| = 0$ for $1 \leq i \leq n$. Making the change of basis

$$e'_1 = e_1, \quad e'_i = e_i + \alpha_{1,i} e_1, \quad 2 \leq i \leq n - 1$$

the multiplication of A simply becomes

$$(*) \quad e_1 e_1 = e_n, \quad e_1 e_i = e_i e_1 = e_i e_i = 0, \quad e_i e_j = \alpha_{ij} e_n, \quad 2 \leq i, j \leq n - 1.$$

We consider now the subalgebra $M : \{e_2, \dots, e_n\}$. Note that M cannot be abelian, since otherwise $(*)$ becomes an algebra λ_2 , which is level one. Thus, we have the following two subcases, which will complete the case $\dim(A^2) = 1$.

Case 1.1.1. Assume that M is Lie. Since M is also not abelian, we are free to choose $\alpha_{2,3} = 1, \alpha_{3,2} = -1$. Taking the degeneration

$$\begin{cases} g_t(e_1) = t^{-2}e_n, & g_t(e_2) = 2t^{-1}e_2 - t^{-2}e_n, \\ g_t(e_3) = t^{-1}e_1 - t^{-2}e_n, & g_t(e_n) = t^{-2}e_3, \\ g_t(e_i) = t^{-2}e_i, & 4 \leq i \leq n - 1, \end{cases}$$

we obtain that A degenerates to $L_4(\frac{1}{4})$.

Case 1.1.2. Assume that M is non-Lie. Then we may assume $e_2e_2 \neq 0$, moreover $e_2e_2 = e_n$. Taking the degeneration

$$g_t(e_1) = e_1, \quad g_t(e_2) = e_2, \quad g_t(e_n) = e_n, \quad g_t(e_i) = t^{-1}e_i, \quad 3 \leq i \leq n - 1,$$

we obtain that A degenerates to L_5 .

Case 1.2. We now assume that $\dim(A^2) \geq 2$. Let $A = \{e_1, \dots, e_n\}$, and $A^2 = \{e_{k+1}, \dots, e_n\}$. We consider five logically exhaustive cases in which the products $e_i e_1, e_1 e_i$ fall in the square A^2 in different ways. It is obvious that we may always assume $e_1 e_1 = e_{k+1}$.

Case 1.2.1. Assume $e_i e_1 \in \text{span}\{e_{k+1}\}$ for $1 \leq i \leq k$, and let $e_1 e_2 \notin \text{span}\{e_{k+1}\}$, so that we have the multiplication:

$$e_1 e_1 = e_{k+1}, \quad e_1 e_2 = e_{k+2}, \quad e_i e_1 = \gamma_{i,1}^{k+1} e_{k+1}, \quad 2 \leq i \leq k,$$

$$e_i e_j = \sum_{\ell=k+1}^n \gamma_{i,j}^\ell e_\ell, \quad 2 \leq i, j \leq k.$$

Applying Lemma 1 for the elements $(\gamma_{1,1}^{k+2}, \gamma_{1,2}^{k+2}, \gamma_{2,1}^{k+2}, \gamma_{2,2}^{k+2}) = (0, 0, 1, \gamma_{2,2}^{k+2})$ we obtain that A degenerates to $L_4(\alpha)$ or L_5 .

Case 1.2.2. Let there exist i such that $e_i e_1 \notin \text{span}\{e_{k+1}\}$. Without loss of generality, we may put $e_2 e_1 = e_{k+2}$, so that our products are

$$e_1 e_1 = e_{k+1}, \quad e_2 e_1 = e_{k+2}, \quad e_i e_1 = \sum_{\ell=k+1}^n \gamma_{i,1}^\ell e_\ell, \quad 3 \leq i \leq k,$$

$$e_1 e_i = \sum_{\ell=k+1}^n \gamma_{1,i}^\ell e_\ell, \quad 2 \leq i \leq k, \quad e_i e_j = \sum_{\ell=k+1}^n \gamma_{i,j}^\ell e_\ell, \quad 2 \leq i, j \leq k.$$

If $\gamma_{1,2}^{k+2} \neq -1$, then applying Lemma 1 for the $(\gamma_{1,1}^{k+2}, \gamma_{1,2}^{k+2}, \gamma_{2,1}^{k+2}, \gamma_{2,2}^{k+2}) = (0, \gamma_{1,2}^{k+2}, 1, \gamma_{2,2}^{k+2})$ we obtain that L degenerates to $L_4(\alpha)$ or L_5 .

If $\gamma_{1,2}^{k+2} = -1$, then taking the change of basis

$$e'_2 = e_2 + \eta e_1, \quad e'_{k+2} = e_{k+2} + \eta e_{k+1}, \quad e'_i = e_i, \quad 1 \leq i (i \neq 2, k+2) \leq n,$$

we obtain that

$$e'_1 e'_1 = e'_{k+1}, \quad e'_2 e'_1 = e'_{k+2}, \quad e'_1 e'_2 = (2\eta + \gamma_{1,2}^{k+1})e'_{k+1} - e'_{k+2} + \sum_{\ell=k+3}^n \gamma_{1,i}^\ell e_\ell,$$

$$e'_i e'_1 = \sum_{\ell=k+1}^n \gamma_{i,1}^\ell e'_\ell, \quad e'_1 e'_i = \sum_{\ell=k+1}^n \gamma_{1,i}^\ell e'_\ell, \quad 3 \leq i \leq k,$$

$$e'_i e'_j = \sum_{\ell=k+1}^n \gamma_{i,j}^\ell e'_\ell \quad 2 \leq i, j \leq k.$$

Taking the value of η such that $\gamma_{1,2}^{k+1'} = 2\eta + \gamma_{1,2}^{k+1} \neq 0$ we apply the Lemma 1 for the elements $(\gamma_{1,1}^{k+1'}, \gamma_{1,2}^{k+1'}, \gamma_{2,1}^{k+1'}, \gamma_{2,2}^{k+1'}) = (1, \gamma_{1,2}^{k+1'}, 0, \gamma_{2,2}^{k+1'})$ and obtain that A degenerates to $L_4(\alpha)$ or L_5 .

Case 1.2.3. Now suppose that $e_i e_1, e_1 e_i \in \text{span}\{e_{k+1}\}$ for $1 \leq i \leq k$ and there exist j such that $e_j e_j \notin \text{span}\{e_{k+1}\}$. In this case without loss of generality, we may suppose $e_2 e_2 = e_{k+2}$, so that our multiplication becomes

$$e_1 e_1 = e_{k+1}, \quad e_2 e_2 = e_{k+2}, \quad e_i e_1 = \gamma_{i,1}^{k+1} e_{k+1}, \quad e_1 e_i = \gamma_{1,i}^{k+1} e_{k+1}, \quad 2 \leq i \leq k,$$

$$e_i e_j = \sum_{\ell=k+1}^n \gamma_{ij}^\ell e_\ell, \quad 3 \leq i, j \leq k.$$

If $(\gamma_{1,2}^{k+1}, \gamma_{2,1}^{k+1}) \neq (0, 0)$ then applying Lemma 1 for the $(\gamma_{1,1}^{k+1}, \gamma_{1,2}^{k+1}, \gamma_{2,1}^{k+1}, \gamma_{2,2}^{k+1}) = (1, \gamma_{1,2}^{k+1}, \gamma_{2,1}^{k+1}, 0)$ we obtain that A degenerates to $L_4(\alpha)$ or L_5 .

If $\gamma_{1,2}^{k+1} = \gamma_{2,1}^{k+1} = 0$, then taking the degeneration

$$g_t(e_1) = t^{-1}e_1, \quad g_t(e_2) = t^{-1}e_2, \quad g_t(e_i) = t^{-2}e_i, \quad 3 \leq i \leq n,$$

we degenerate to the algebra

$$\lambda_2 \oplus \lambda_2 : e_1 e_1 = e_{k+1}, \quad e_2 e_2 = e_{k+2}.$$

By Example 1, we obtain that algebra $\lambda_2 \oplus \lambda_2$ degenerates to the algebra L_5 .

Case 1.2.4. Now we suppose that $e_i e_i, e_1 e_j, e_i e_1 \in \text{span}\{e_{k+1}\}$ for $1 \leq i \leq k$. Let there exist i, j ($2 \leq i, j \leq k$) such that $e_i e_j \notin \text{span}\{e_{k+1}\}$. Without loss of generality we can suppose $i = 2, j = 3$ moreover, if $e_2 e_3 + e_3 e_2 \neq 0$, then taking $e'_2 = e_2 + e_3$, we get that $e'_2 e'_2 = e_2 e_2 + e_2 e_3 + e_3 e_2 + e_3 e_3 \notin \{e_{k+1}\}$ which have the situation of Case 1.2.3. Therefore, we may suppose

$$e_1 e_1 = e_{k+1}, \quad e_1 e_2 = \alpha_{1,2} e_{k+1}, \quad e_2 e_1 = \alpha_{2,1} e_{k+1}, \quad e_2 e_2 = \alpha_{2,2} e_{k+1},$$

$$e_1 e_3 = \alpha_{1,3} e_{k+1}, \quad e_3 e_1 = \alpha_{3,1} e_{k+1}, \quad e_3 e_3 = \alpha_{3,3} e_{k+1},$$

$$e_2 e_3 = e_{k+2}, \quad e_3 e_2 = -e_{k+2}.$$

Now applying the degeneration

$$\begin{cases} g_t(e_1) = t^{-2}e_1, & g_t(e_2) = t^{-3}e_2, \\ g_t(e_3) = t^{-3}e_3, & g_t(e_{k+1}) = t^{-4}e_{k+1}, \\ g_t(e_i) = t^{-5}e_i, & 4 \leq i \leq n. \end{cases}$$

we obtain the products

$$e_1 e_1 = e_{k+1}, \quad e_2 e_3 = e_{k+2}, \quad e_3 e_2 = -e_{k+2}.$$

which by Example 1 degenerates to the algebra L_5 .

Case 1.2.5. Now we suppose that $e_i e_j \in \text{span}\{e_{k+1}\}$ for all i, j ($1 \leq i, j \leq k$). Thus we have

$$e_1 e_1 = e_{k+1}, \quad e_i e_j = \alpha_{i,j} e_{k+1}.$$

Since $\dim(A^2) \geq 2$, then we have that there exist i ($2 \leq i \leq k$), such that $e_i e_{k+1}$ or $e_{k+1} e_i$ is non zero, which we can suppose as e_{k+2} .

Let $i = 1$, then we have $e_1 e_1 = e_{k+1}, e_1 e_{k+1} = e_{k+2}$.

If $e_{k+1} e_1 + e_1 e_{k+1} \neq 0$, then using the Lemma 1 for the basis elements $\{e_1, e_{k+1}, e_{k+2}\}$ we have that A degenerates to the algebra $L_4(\alpha)$ or L_5 .

If $e_{k+1} e_1 = -e_1 e_{k+1} = -e_{k+2}$ then making the change $e'_{k+1} = e_{k+1} - e_{k+2}$, we have that

$$e_1 e_1 = e_{k+1} + e_{k+2}, \quad e_1 e_{k+1} = e_{k+2}, \quad e_{k+1} e_1 = -e_{k+2}.$$

Again applying Lemma 1 for the basis elements $\{e_1, e_{k+1}, e_{k+2}\}$, i.e. $(\gamma_{1,1}^{k+2}, \gamma_{1,k+1}^{k+2}, \gamma_{k+1,1}^{k+2}, \gamma_{k+1,k+1}^{k+2}) = (1, 1, -1, \gamma_{k+1,k+1}^{k+2})$ we have that A degenerates to the algebra $L_4(\alpha)$ or L_5 .

Let $i \neq 1$, then we can suppose $i = 2$ and we have $e_1 e_1 = e_{k+1}, e_2 e_{k+1} = e_{k+2}$.

Similarly to the case $i = 1$ if $e_{k+1} e_2 + e_2 e_{k+1} \neq 0$, then using Lemma 1 for the basis elements $\{e_2, e_{k+1}, e_{k+2}\}$ we have that A degenerates to the algebra $L_4(\alpha)$ or L_5 .

If $e_{k+1} e_2 = -e_2 e_{k+1} = -e_{k+2}$ then making the change $e'_{k+1} = e_{k+1} - e_{k+2}$, we have that

$$e_1 e_1 = e_{k+1} + e_{k+2}, \quad e_1 e_{k+1} = e_{k+2}, \quad e_{k+1} e_1 = -e_{k+2}.$$

Again applying Lemma 1 for the basis elements $\{e_1, e_{k+1}, e_{k+2}\}$, i.e. $(\gamma_{1,1}^{k+2}, \gamma_{1,k+1}^{k+2}, \gamma_{k+1,1}^{k+2}, \gamma_{k+1,k+1}^{k+2}) = (1, 1, -1, \gamma_{k+1,k+1}^{k+2})$ we have that A degenerates to the algebra $L_4(\alpha)$ or L_5 .

Case 2. Let $A \in \text{Nil}_n(\mathbb{C})$ be antisymmetric. It should be noted that if $\dim(A^2) = 1$, then we have that $A^3 = 0$. Thus, A is a Lie algebra. In [12] it is shown that any nilpotent Lie algebra with condition $\dim(A^2) = 1, A^3 = 0$ degenerates to algebra $n_{5,1}$.

Therefore, we consider case $\dim(A^2) \geq 2$. Assume that $\{e_1, e_2, \dots, e_k, e_{k+1}, \dots, e_n\}$ be a basis of A , and $\{e_{k+1}, e_{k+2}, \dots, e_n\}$ be a basis of A^2 .

Then, without loss of generality, we can assume $e_1 e_2 = e_{k+1}, e_2 e_1 = -e_{k+1}$.

Below, we show that it may always be assumed

$$e_1 e_2 = e_4, \quad e_1 e_3 = e_5.$$

- Let there exists i_0 such that $e_1e_{i_0} \notin \text{span}\langle x_{k+1} \rangle$. Then taking

$$e'_1 = e_1, e'_2 = e_2, e'_3 = e_{i_0}, e'_4 = e_{k+1}, e'_5 = e_1e_{i_0}$$

we obtain $e'_1e'_2 = e'_4, e'_1e'_3 = e'_5$.

- Let $e_1e_i \in \text{span}\{e_{k+1}\}$ for all $3 \leq i \leq k$ and there exists some i_0 such that $e_2e_{i_0} \notin \text{span}\langle x_{k+1} \rangle$. According to symmetrically of e_1 and e_2 , similarly to the previous case we can choose a basis $\{e'_1, e'_2, \dots, e'_n\}$ with condition $e'_1e'_2 = e'_4, e'_1e'_3 = e'_5$.
- Let $x_1x_i, x_2x_i \in \text{span}\{e_{k+1}\}$ for all $3 \leq i \leq k$. We set $e_1e_i = \alpha_i e_{k+1}$ and $e_2e_i = \beta_i x_{k+1}$. Let e_{i_0} and e_{j_0} be generators of A such that $e_{i_0}e_{j_0} \notin \text{span}\{e_{k+1}\}$. Since $\dim(A^2) \geq 2$ one can assume $e_{i_0}e_{j_0} = e_{k+2}$.

Putting

$$e'_1 = e_1 + Ae_{i_0}, e'_2 = e_2, e'_3 = e_{j_0}, e'_4 = (1 - A\beta_{i_0})e_{k+1}, e'_5 = Ae_{k+2} + \alpha_{i_0}e_{k+1}$$

with $A(1 - A\beta_{i_0}) \neq 0$, we deduce $e'_1e'_2 = e'_4, e'_1e'_3 = e'_5$.

- Let $e_i e_j \in \text{span}\{e_{k+1}\}$ for all $1 \leq i, j \leq k$. Then for some i_0 we have $e_{i_0}e_{k+1} \neq 0$. Without loss of generality, one can assume $e_1e_{k+1} = e_{k+2}$.

– If $k \geq 3$, then setting

$$e'_1 = e_1, e'_2 = e_2, e'_3 = e_3 + e_{k+1}, e'_4 = e_{k+1}, e'_5 = e_{k+2} + \alpha_{1,3}e_{k+1},$$

we obtain $e'_1e'_2 = e'_4, e'_1e'_3 = e'_5$.

– If $k = 2$, then we have $e_1e_2 = e_3, e_1e_3 = e_4$. It is not difficult to obtain that $e_1e_4 = e_5$ or $e_2e_3 = e_5$ (because of $n \geq 5$). Indeed, taking

$$e'_1 = e_1, e'_2 = e_2, e'_3 = e_4, e'_4 = e_3, e'_5 = e_5$$

in the case of $e_1e_4 = e_5$ and

$$e'_1 = -e_3, e'_2 = e_1, e'_3 = e_4, e'_4 = e_2, e'_5 = e_5$$

in the case of $e_2e_3 = e_5$, we derive the products $e_1e_2 = e_4, e_1e_3 = e_5$.

Thus, there exists a basis $\{e_1, e_2, e_3, \dots, e_n\}$ of A with the products

$$e_1e_2 = e_4, e_1e_3 = e_5.$$

Note that A degenerates to the algebra with multiplication:

$$e_1e_2 = e_4, e_1e_3 = e_5, e_2e_3 = \gamma_4x_4 + \gamma_4x_5$$

via the following degeneration:

$$g_t : \begin{cases} g_t(e_1) = t^{-2}e_1, & g_t(e_2) = t^{-2}e_2, & g_t(e_3) = t^{-2}e_3, \\ g_t(e_4) = t^{-4}e_4, & g_t(e_5) = t^{-4}e_5, & g_t(e_i) = t^{-3}e_i, \end{cases} \quad 6 \leq i \leq n.$$

From the change of basis $e'_2 = e_2 - \gamma_5e_1, e'_3 = e_3 + \gamma_4e_1$, we obtain that this algebra is isomorphic to $n_{5,2} \oplus \text{ab}_{n-5}$. \square

Acknowledgments

The authors are very grateful to the reviewer for his/her valuable comments and suggestions which helped to improve the presentation. The work was supported by National Science Foundation (United States) award number 1658672.

References

- [1] J.R. Nielsen (Ed.), Niels Bohr Collected Works, 3: The Correspondence Principle, North-Holland Publishing, Amsterdam, 1976, pp. 1918–1923.
- [2] M. Gerstenhaber, On the deformations of rings and algebras, *Ann. of Math.* 79 (1964) 59–103.
- [3] F. Bayen, M. Flato, C. Fronsdal, A. Lichnerowicz, D. Sternheimer, Deformation theory and quantization. I. Deformations of symplectic structures, *Ann. Phys.* 111 (1978) 61–110.
- [4] S. Waldmann, Poisson-geometrie und deformationsquantisierung, in: *Eine Einführung*, Springer, 2007.
- [5] K. Costello, O. Gwilliam, Factorization algebras in perturbative quantum field theory, 2016, p. 318.
- [6] Dütsch, K. Fredenhagen, Perturbative algebraic quantum field theory and deformation quantization, in: *Proceedings of the Conference on Mathematical Physics in Mathematics and Physics*. Siena June 20–25 (2000).
- [7] D. Hilbert, Mathematical problems, *Bull. Amer. Math. Soc.* 8 (10) (1902) 437–479.
- [8] K. Kodaira, D.C. Spencer, On deformations of complex analytic structures i-ii, *Ann. of Math.* 67 (2) (1958) 328–466.
- [9] W. Lowen, M. Van Den Bergh, Deformation theory of Abelian categories, *Trans. Amer. Math. Soc.* 358 (12) (2006) 5441–5483.
- [10] E. İnönü, E.P. Wigner, On the contraction of groups and their representations, *Proc. Natl. Acad. Sci.* 39 (6) (1953) 510–524.
- [11] A.Kh. Khudoyberdiyev, B.A. Omirov, The classification of algebras of level one, *Linear Algebra Appl.* 439 (11) (2013) 3460–3463.
- [12] A.Kh. Khudoyberdiyev, The classification of Algebras of level two, *J. Geom. Phys.* 98 (2015) 13–20.
- [13] J.-L. Loday, Une version non commutative des algèbres de Leibniz, *Enseign. Math.* 39 (1993) 269–293.
- [14] S. Albeverio, B.A. Omirov, I.S. Rakhimov, Varieties of nilpotent complex Leibniz algebras of dimensions less than five, *Comm. Algebra* 33 (5) (2015) 1575–1585.

- [15] D. Belțiță, B. Cahen, Contractions of Lie algebras with 2-dimensional generic coadjoint orbits, *Linear Algebra Appl.* 466 (2015) 41–63.
- [16] D. Burde, K. Dekimpe, K. Vercammen, Affine actions on Lie groups and post-Lie algebra structures, *Linear Algebra Appl.* 437 (5) (2012) 1250–1263.
- [17] V.V. Gorbatsevich, Contractions and degenerations of finite-dimensional algebras, *Sov. Math. (Iz. VUZ)*. 35 (10) (1991) 17–24.
- [18] F. Grunewald, J. O'Halloran, Varieties of nilpotent Lie algebras of dimension less than six, *J. Algebra* 112 (2) (1988) 315–325.
- [19] J.M. Casas, A.Kh. Khudoyberdiyev, M. Ladra, B.A. Omirov, On the degenerations of solvable Leibniz algebras, *Linear Algebra Appl.* 439 (2) (2013) 472–487.
- [20] A.Kh. Khudoyberdiyev, B.A. Omirov, Infinitesimal deformations of null-filiform Leibniz superalgebras, *J. Geom. Phys.* 74 (2013) 370–380.
- [21] S. Albeverio, B.A. Omirov, I.S. Rakhimov, Classification of 4-dimensional nilpotent complex Leibniz algebras, *Extracta Math.* 21 (3) (2006) 197–210.